



PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : C12N 15/11, A61K 31/70, C07H 21/04	A1	(11) International Publication Number: WO 96/32474 (43) International Publication Date: 17 October 1996 (17.10.96)
(21) International Application Number: PCT/US96/04605 (22) International Filing Date: 4 April 1996 (04.04.96) (30) Priority Data: 08/420,672 12 April 1995 (12.04.95) US (71) Applicant: HYBRIDON, INC. [US/US]; One Innovation Drive, Worcester, MA 01605 (US). (72) Inventors: KANDIMALLA, Ekambar, R.; Apartment #305, 286 Plantation Street, Worcester, MA 01604 (US). AGRAWAL, Sudhir, 61 Lamplighter Drive, Shrewsbury, MA 01545 (US). (74) Agents: KERNER, Ann-Louise; Lappin & Kusmer, Two Hundred State Street, Boston, MA 02109 (US) et al.		(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>
(54) Title: COOPERATIVE OLIGONUCLEOTIDES (57) Abstract Disclosed is a composition comprising at least two synthetic, cooperative oligonucleotides, each comprising a region complementary to one of tandem, non-overlapping regions of a target single-stranded nucleic acid, and each further comprising a dimerization domain at a terminus of each of the oligonucleotides, the dimerization domains of the oligonucleotides being complementary to each other. Also disclosed are duplex structures, ternary complexes, pharmaceutical formulations, and methods utilizing the cooperative oligonucleotides of the invention.		

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LI	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

COOPERATIVE OLIGONUCLEOTIDESCROSS-REFERENCE TO RELATED APPLICATIONS

- 5 This application is a continuation-in-part of
copending Patent Application Serial No. 08/420,672
entitled "COOPERATIVE OLIGONUCLEOTIDES", filed April 12,
1995.

10 BACKGROUND OF THE INVENTION

- Progress in chemical synthesis of nuclease resistant
oligonucleotides (*Methods Mol. Biol.* (1993) Vol. 20, (Agrawal,
ed.) Humana Press, Totowa, NJ) and developments in large
15 scale solid phase synthesis of oligonucleotides
((Agrawal, ed.) *Methods Mol. Biol.* (1993) Vol. 20, Humana
Press, Totowa, NJ); Padmapriya et al. (1994) *Antisense Res.*
Dev. 4:185-199) has permitted antisense oligonucleotides
to advance to human clinical trials (Bayever et al.
20 (1993) *Antisense Res. Dev.* 3:383-390). In principle, antisense
oligonucleotides utilize highly sequence-specific
complementary nucleo-base recognition of target nucleic
acids through Watson-Crick hydrogen bonding between A and
T, and G and C, that leads to the development of less
25 toxic and more site specific chemotherapeutic agents
(Stephenson et al. (1978) *Proc. Natl. Acad. Sci. (USA)* 75:285-
288). As per theoretical calculations, an
oligonucleotide of 13 or more bases long should bind to a
unique sequence that occurs only once in a eucaryotic
30 mRNA pool.

- Contrary to the popular belief, it was recently
shown that the increase in the length of an antisense
oligonucleotide beyond the minimum length that can
35 hybridize to the target (i.e. 11-14 bases) decreases its
specificity rather than increasing (Woolf et al. (1992)

-2-

Proc. Natl. Acad. Sci. (USA) 89:7305-7309). Potentially, this decrease in hybridization specificity would lead to non-sequence-specific target binding and subsequent increased toxicity (Stein et al. (1993) *Science* 261:1004-1012).

5

Thus, what is needed are improved antisense oligonucleotides optimized for therapeutic and diagnostic use which have improved affinity, specificity, and biological activity, and little or no toxicity.

10

SUMMARY OF THE INVENTION

The present invention provides cooperative oligonucleotides with improved sequence specificity for a single-stranded target, reduced toxicity, and improved biological activity as antisense molecules.

Surprisingly, it has been discovered that two short oligonucleotides (25 nucleotides or less) bind to adjacent sites on the target nucleic acid in a cooperative manner, allowing for an interaction with greater sequence specificity than can a single longer oligonucleotide having a length equal to the two shorter oligonucleotides.

25

Accordingly, in a first aspect, the present invention provides a composition including at least two synthetic cooperative oligonucleotides, each comprising a region complementary to one of tandem, non-overlapping regions of a target single-stranded nucleic acid, and a dimerization domain at a terminus of each of the oligonucleotides. The dimerization domains of the cooperative oligonucleotides are complementary to each other, and the target nucleic acid being an mRNA, single-stranded viral DNA, or single-stranded viral RNA.

35

-3-

In some preferred embodiments, the oligonucleotides each are complementary to tandem regions of the target nucleic acid that are separated by 0 to 3 bases. In some preferred embodiments, each of the oligonucleotides are
5 about 9 to 25 nucleotides in length.

In one embodiment, the composition consists of two cooperative oligonucleotides, the dimerization domain of a first or one of the oligonucleotides being located at
10 its 3' terminal portion, and being complementary to the dimerization domain of a second or the other oligonucleotide which is located at its 5' terminal portion. Alternatively, the dimerization domain of the first cooperative oligonucleotide is located at its 3'
15 terminal portion, and is complementary to the dimerization domain of a second oligonucleotide which is located at its 3' terminal portion. Alternatively, the dimerization domain of the first cooperative oligonucleotide is located at its 5' terminal portion,
20 and is complementary to a dimerization domain of the second oligonucleotide which is located at its 5' terminal portion.

The invention provides in another aspect a duplex
25 structure comprising first and second synthetic cooperative oligonucleotides, each oligonucleotide comprising a region complementary to the non-overlapping, tandem regions of the target nucleic acid which is an mRNA, single-stranded viral RNA, or single-stranded viral
30 DNA. The first oligonucleotide in the duplex has a terminal dimerization domain complementary and hybridized to the dimerization domain of the second oligonucleotide. In some embodiments, each of the oligonucleotides are about 9 to 25 nucleotides in length, and in others, the
35 dimerization domains of the first and second oligonucleotides each comprise about 3 to 7 nucleotides. In some embodiments, the invention provides first and

-4-

second oligonucleotides which are complementary to tandem regions of the target nucleic acid separated by 0 to 3 bases. In another embodiment, the duplex structure is hybridized to the target nucleic acid.

5

The invention also provides pharmaceutical formulations containing the compositions or duplex structures described above, and methods of inhibiting the expression of a nucleic acid *in vitro* comprising the step of
10 treating the nucleic acid with the pharmaceutical formulations of the invention. In some embodiments, the first and second oligonucleotides are complementary to an HIV DNA or an HIV RNA.

15 In another aspect, the invention provides a ternary structure comprising a first synthetic cooperative oligonucleotide, a second synthetic cooperative oligonucleotide, and a third synthetic cooperative oligonucleotide, each oligonucleotide comprising a region
20 complementary to one of tandem, non-overlapping regions of a target nucleic acid, and each comprising a dimerization domain at one or both of their termini. The dimerization domain of the first oligonucleotide is complementary and hybridized to a first dimerization
25 domain at one terminus of the third oligonucleotide and the dimerization domain of the second oligonucleotide is complementary and hybridized to a second dimerization domain at the other terminus of the third oligonucleotide when the first, second, and third oligonucleotides are
30 hybridized to the target nucleic acid. The target nucleic acid is an mRNA, a single-stranded viral RNA, or a single-stranded viral DNA. In some embodiments, the ternary structure is hybridized to the target nucleic acid.

35

-5-

In another embodiment, the invention provides methods of inhibiting the expression of a nucleic acid comprising the step of contacting the nucleic acid with at least one cooperative oligonucleotide of the

5 invention.

BRIEF DESCRIPTION OF THE DRAWINGS

The foregoing and other objects of the present invention, the various features thereof, as well as the invention itself may be more fully understood from the following description, when read together with the accompanying drawings in which:

FIG. 1A is a schematic representation of the cooperative binding of two short oligonucleotides to tandem sites;

FIG. 1B is a schematic representation of the binding to adjacent sites on a target nucleic acid of cooperative oligonucleotides that have extended antisense dimerization domains and their dimerization;

FIG. 1C is a schematic representation of the binding of three cooperative oligonucleotides of the invention to adjacent sites on a target nucleic acid;

FIG. 2A is a graphic representation showing the thermal melting profile (dA/dT vs. T) of oligonucleotides 1-7 shown in Table 2 with their DNA target;

FIG. 2B is a graphic representation showing the thermal melting profile (dA/dT vs. T) of oligonucleotides 1+2, 1+3, 1+4, and 5 shown in Table 2 with their DNA target;

FIG. 3 is a graphic representation showing the thermal melting profiles (dA/dT vs. T) of the oligonucleotide combinations with extended antisense dimerization domains (10+14, 11+15, 9+14, 12+16, and 13+17);

-7-

FIG. 4A is an autoradiogram showing the RNase H hydrolysis pattern of the RNA target sequence in the presence of oligonucleotides 5, 1, 2, 1+2, 14, 10, and 10+14 at different time points;

5

FIG. 4B is an autoradiogram showing the RNase H hydrolysis pattern of the RNA target sequence in the presence of oligonucleotides 5, 13, 17, and 13+17 at different time points;

10

FIG. 5 is an autoradiogram showing the RNase H hydrolysis pattern of RNA target in the presence of the mismatched oligonucleotides 23, 24, 18 and 19 compared to the control matched oligonucleotide 5 and 1 at different time points;

15

FIG. 6 is a graphic representation showing the ability of cooperative oligonucleotide oligonucleotides 1+2 (--◇--), and 13+17 (--O--), and control oligonucleotides 5 (--□--) and 20 (--Δ--) at varying concentrations to inhibit HIV-1 in a cell culture system;

20

FIG. 7 is a graphic representation showing the percent inhibition of HIV-1 in cell cultures by cooperative antisense oligonucleotides 1+2, 13+17, 9+14, 10+14, and 12+16 and by control antisense oligonucleotides 5 and 20, present at two different concentrations; and

30

FIG. 8 is a graphic representation showing the relationship between meeting temperature (T_m) and percent HIV-1 inhibition for cooperative oligonucleotides 10+14, 12+16, and 13+17.

35

DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

Cooperative interactions between biological macromolecules are important in nature. For example, the cooperative interactions between proteins and nucleic acids are vital for the regulation of gene expression. Cooperative interactions serve to improve sequence specificity, affinity, and biological activity (Ptashne (1986) *A Genetic Switch*; Blackwell Scientific Publications and Cell Press: Palo Alto, CA). Cooperative binding of drugs to DNA (Asseline et al. (1984) *Proc. Natl. Acad. Sci. (USA)* 81:3297-3301; Rao et al. (1991) *J. Org. Chem.* 56:786-797), of oligonucleotides or their conjugates to single stranded DNA (Tazawa et al. (1972) *J. Mol. Biol.* 66:115-130; Maher et al. (1988) *Nucl. Acids Res.* 16:3341-3358; Springgate et al. (1973) *Biopolymers* 12:2241-2260; and Gryaznov et al. (1993) *Nucl. Acids Res.* 21:5909-5915), of oligonucleotides to RNA (Maher III et al. (1987) *Arch. Biochem. Biophys.* 253:214-220), and of oligonucleotides to double-stranded DNA through triplex formation (Strobel et al. (1989) *J. Am. Chem. Soc.* 111:7286-7287; Distefano et al. (1991) *J. Am. Chem. Soc.* 113:5901-5902; Distefano et al. (1992) *J. Am. Chem. Soc.* 114:11006-11007; Colocci et al. (1993) *J. Am. Chem. Soc.* 115:4468-4473; Colocci et al. (1994) *J. Am. Chem. Soc.* 116:785-786) has been documented. Although these studies demonstrated the advantages of using cooperative interactions for small molecule-based drug development, there are no reports of optimizing the design of cooperative oligonucleotides for therapeutic uses.

30

The present invention provides synthetic oligonucleotides which interact with mRNA, single-stranded viral RNA, or single-stranded viral DNA ("target nucleic acids"), and have improved affinity, specificity, and biological activity as antisense molecules. At least

35

-9-

two of the oligonucleotides of the invention are used to interact with a target nucleic acid, thereby enabling them to interact cooperatively, synergistically enhancing their ability (singly) to inhibit expression of the target nucleic acid.

The term "synthetic oligonucleotide" for purposes of this invention includes chemically synthesized polymers of about 7 to about 25, and preferably from about 9 to about 23 nucleotide monomers (nucleotide bases) connected together or linked by at least one 5' to 3' internucleotide linkage.

Some cooperative oligonucleotides of the invention are complementary to non-overlapping, tandem regions of the target nucleic acid, as shown in FIG. 1A, while others are complementary to adjacent sites (FIGS. 1B and 1C). At least two of these oligonucleotides can be used to control target nucleic acid expression.

For purposes of the invention, the term "oligonucleotide complementary to a target nucleic acid" is intended to mean an oligonucleotide sequence that binds to the nucleic acid sequence under physiological conditions, e.g., by Watson-Crick base pairing (interaction between oligonucleotide and single-stranded nucleic acid in antiparallel orientation) or by Hoogsteen base pairing (interaction between oligonucleotide and double-stranded or single-stranded nucleic acid in parallel orientation) or by any other means including in the case of a oligonucleotide binding to RNA, pseudoknot formation. Such binding (by Watson-Crick base pairing) under physiological conditions is measured as a practical matter by observing interference with the function of the nucleic acid sequence.

The inhibitory ability of the cooperative oligonucleotides of the invention is enhanced even

-10-

further when these oligonucleotides also include a terminal portion (i.e., a "dimerization domain") which is not complementary to the target nucleic acid, but rather which is complementary to each other, thereby enabling the formation of a dimers (FIG. 1B). The interaction of these cooperative oligonucleotides with the target nucleic acid leads to the formation of a more stable ternary complex as the result of dimerization of the complementary dimerization domains of these oligonucleotides. When the cooperative oligonucleotides of the invention have dimerization domains and hybridize together to form a duplex, the regions of the cooperative oligonucleotides which are complementary to the target nucleic acid may be separated by 0 to 3 bases.

The cooperative oligonucleotides of the invention may have any nucleotide sequence, as long as a portion of its sequence is complementary to a portion of a target nucleic acid. Any nucleic acid may be targeted by the cooperative oligonucleotides of the invention including viral, bacterial, and cellular genes, mRNAs, or cDNAs. Further, the terminal dimerization domains of cooperative oligonucleotides which form duplexes with each other may not be not complementary to the target nucleic acid. These dimerization domains may be at the 3' termini of both cooperative oligonucleotides, at the 5' termini of both cooperative oligonucleotides, or at the 3' terminus of one cooperative oligonucleotide and the 5' terminus of the other cooperative oligonucleotide.

The cooperative oligonucleotides of the invention are composed of deoxyribonucleotides, ribonucleotides, or any combination thereof, with the 5' end of one nucleotide and the 3' end of another nucleotide being covalently linked, in some cases, via a phosphodiester internucleotide linkage. The oligonucleotides can be prepared by art recognized methods such as

-11-

- phosphoramidate, H-phosphonate chemistry, or methylphosphoramidate chemistry (*see, e.g.,* Uhlmann et al. (1990) *Chem. Rev.* 90:543-584; Agrawal et al. (1987) *Tetrahedron. Lett.* 28:(31):3539-3542; Caruthers et al. (1987) *Meth. Enzymol.* 154:287-313; U.S. Patent 5,149,798) which can be carried out manually or by an automated synthesizer and then processed (reviewed in Agrawal et al. (1992) *Trends Biotechnol.* 10:152-158).
- 10 The oligonucleotides of the invention may also be modified in a number of ways, for example, to enhance stability, without compromising their ability to hybridize to nucleotide sequences contained within a targeted region of a particular gene.
- 15 The term "modified oligonucleotide" as used herein describes an oligonucleotide in which at least two of its nucleotides are covalently linked via a synthetic linkage, i.e., a linkage other than a phosphodiester linkage between the 5' end of one nucleotide and the 3' end of another nucleotide in which the 5' nucleotide phosphate has been replaced with any number of chemical groups. For example, oligonucleotides with phosphorothioate linkages can be prepared using methods well known in the field such as methoxyphosphoramidite (*see, e.g.,* Agrawal et al. (1988) *Proc. Natl. Acad. Sci. (USA)* 85:7079-7083) or H-phosphonate (*see, e.g.,* Froehler (1986) *Tetrahedron Lett.* 27:5575-5578) chemistry. The synthetic methods described in Bergot et al. (*J. Chromatog.* (1992) 559:35-42) can also be used. Examples of other chemical groups include alkylphosphonates, phosphorodithioates, alkylphosphonothioates, phosphoramidates, phosphoramidites, phosphate esters, 2-O-methyls, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters, and thiono triesters. Any of these chemical groups or linkages may
- 20
- 25
- 30
- 35

-12-

also be substituted with various other chemical groups, e.g., an aminoalkylphosphonate. Oligonucleotides with these chemical groups can be prepared according to known methods (see, e.g., Agrawal and Goodchild (*Tetrahedron Lett.* 5 (1987) 28:3539-3542); Agrawal et al. (*Proc. Natl. Acad. Sci. (USA)* (1988) 85:7079-7083); Uhlmann et al. (*Chem. Rev.* (1990) 90:534-583; and Agrawal et al. (*Trends Biotechnol.* (1992) 10:152-158). U.S. Patent Application Ser. No. (47508-559), filed on August 9, 1995 discloses "inverted" 10 chimeric oligonucleotides comprising one or more nonionic oligonucleotide region (e.g. alkylphosphonate and/or phosphoramidate and/or phosphotriester internucleoside linkage) flanked by one or more region of oligonucleotide phosphorothioate. The phosphorothioate linkages may be 15 mixed Rp and Sp enantiomers, or they may be stereoregular or substantially stereoregular in either Rp or Sp form (see Iyer et al. (1995) *Tetrahedron Asymmetry* 6:1051-1054). Thiono triesters can be prepared according to the methods described in U.S. Ser. No. 08/409,169, filed March 23, 20 1993. Oligonucleotides with phosphorothioate linkages can be prepared using methods well known in the field such as phosphoramidite (see, e.g., Agrawal et al. (1988) *Proc. Natl. Acad. Sci. (USA)* 85:7079-7083). or by H-phosphonate (see, eg., Froehler (1986) *Tetrahedron Lett.* 27:5575-5578) 25 chemistry. The synthetic methods described in Bergot et al. (*J. Chromatog.* (1992) 559:35-42) can also be used.

Oligonucleotides which are self-stabilized are also considered to be modified oligonucleotides useful in the 30 methods of the invention (Tang et al. (1993) *Nucleic Acids Res.* 20:2729-2735). These oligonucleotides comprise two regions: a target hybridizing region; and a self-complementary region having an oligonucleotide sequence complementary to a nucleic acid sequence that is 35 within the self-stabilized oligonucleotide.

-13-

Other modifications include those which are internal or are at the end(s) of the oligonucleotide molecule and include additions to the molecule of the internucleoside phosphate linkages, such as cholesteryl or diamine compounds with varying numbers of carbon residues between the amino groups and terminal ribose, deoxyribose and phosphate modifications which cleave, or crosslink to the opposite chains or to associated enzymes or other proteins which bind to the viral genome. Examples of such modified oligonucleotides include oligonucleotides with a modified base and/or sugar such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide having a sugar which, at both its 3' and 5' positions is attached to a chemical group other than a hydroxyl group (at its 3' position) and other than a phosphate group (at its 5' position).

Yet other examples of modifications to sugars include modifications to the 2' position of the ribose moiety which include but are not limited to 2'-O-substituted with an -O- lower alkyl group containing 1-6 saturated or unsaturated carbon atoms, or with an -O-aryl, or allyl group having 2-6 carbon atoms wherein such -O-alkyl, aryl or allyl group may be unsubstituted or may be substituted, (e.g., with halo, hydroxy, trifluoromethyl cyano, nitro acyl acyloxy, alkoxy, carboxy, carbalkoxyl, or amino groups), or with an amino, or halo group. None of these substitutions are intended to exclude the native 2'-hydroxyl group in the case of ribose or 2'-H- in the case of deoxyribose. PCT Publication No. WO 94/02498 discloses traditional hybrid oligonucleotides having regions of 2'-O-substituted ribonucleotides flanking a DNA core region. U.S. Patent Application Serial No. 08/516,454, filed August 17, 1995, discloses an "inverted" hybrid oligonucleotide which includes an oligonucleotide comprising a 2'-O-substituted (or 2' OH, unsubstituted) RNA region which is in between

-14-

two oligodeoxyribonucleotide regions, a structure that "inverted relative to the "traditional" hybrid oligonucleotides.

5 Other modified oligonucleotides are capped with a nuclease resistance-conferring bulky substituent at their 3' and/or 5' end(s) , or have a substitution in one nonbridging oxygen per nucleotide. Such modifications can be at some or all of the internucleoside linkages, as
10 well as at either or both ends of the oligonucleotide and/or in the interior of the molecule (reviewed in Agrawal et al. (1992) *Trends Biotechnol.* 10:152-158).

 To demonstrate the cooperative nature of the
15 oligonucleotides of the invention, oligonucleotides were prepared as described above and tested for their ability to inhibit the expression of a target gene.

 The target chosen was a sequence in the initiation
20 codon region of *gag* mRNA of HIV-1 (SEQ ID NOS:21 and 22) (Agrawal and Tang (1992) *Antisense Res. Dev.* 2:261). A nonlimiting list of some representative *gag* oligonucleotides are shown in TABLE 1.

-15-

TABLE 1

SEQ ID NO:	Sequence* (3' → 5')	Length (# bases)
21	CTAGAAGGAGAGAGATGGGTGCCGAGAG	Target ^b
22	AGAAGGAGAGAGAGTGGGUGCGAGAGCGUCAGUAUUAAGC	Target ^b
25	GGAGGCUAGAAGGAGAGAGAUGGGUGCGAGAGCGU	Target ^b
1	CCCACGCTC	9
2	TTCCTCTCTCTA	12
3	CTTCCTCTCTCT	12
4	TCTTCCTCTCTC	12
5	TTCCTCTCTCTACCCACGCTC	21
6	CTTCCTCTCTCTGCCACGCTC	22
7	TCTTCCTCTCTCCGCCACGCTC	23
8	CTTCCTCTCTCTA	13
9	TTCCTCTCTCTA G G C	15 15
10	CTTCCTCTCTCT G G C	15
11	CTTCCTCTCTCT G G C C	16
12	CTTCCTCTCTCT G G C C G	17
57	CTTCCTCTCTCT	12
19	G G C C G C G	7
14	CCCACGCTC C C G	12
15	CCCACGCTC C C G G G	13

-16-

TABLE 1 (con't)		
SEQ ID NO:	Sequence ^a (3' → 5')	Length (# bases)
16	CCCACGCTC C C G G C	14
17	CCCACGCTC C C G G C G C	16
18	CCCAC <u>T</u> CTC	9
19	CCA <u>A</u> CTCTC	9
20	TCTTCCTCTCTCTACCCACGCTCTC	25
23	TTCCTCTCTCTACCCAC <u>T</u> CTC	21
24	TTCCTCTCTCTACCA <u>A</u> CTCTC	21
26	TCTTCCTCT T C A C C A G	16
27	CTCTACC A G G A T G G T G A T A C G	21
28	CACGCTCTC C T C A C T C	16
29	TCTTCCTCT T C A C C	14

-17-

TABLE 1 (con't)		
SEQ ID NO:	Sequence ^a (3' → 5')	Length (# bases)
30	CTCTACC A G G A T G G T G A	17
31	CACGCTCTC C T C A C	14
32	TCTTCCTCT T C A	12
33	CTCTACC A G G A T G	13
34	CACGCTCTC C T C	12
35	TCTTCCTCT	9
36	CTCTACC	7
37	CACGCTCTC	9

^a

underlined bases represent mismatches

^b sequence is 5' → 3'

Oligonucleotides 1 (SEQ ID NO:1) and 2 (SEQ ID NO:2) are designed to bind to 21 bases of the target nucleic acid at adjacent sites without any base gap between them (see FIG. 1A and TABLE 1). Thus, contact is expected to be maintained through the 3'-end of the oligonucleotide 1 and the 5'-end of the oligonucleotide 2 when these oligonucleotides bind to the target sequence at the adjacent sites. This

-18-

results in cooperativity in the interactions of these two oligonucleotides. Oligonucleotides 3 (SEQ ID NO:3) and 4 (SEQ ID NO:4) bind to the same site as oligonucleotide 2 but are separated by 1 and 2 bases on the target sequence, gaps, respectively, from the binding site of oligonucleotide 1. Because of this gap these oligonucleotides are expected not to show any cooperativity in the binding of these oligonucleotide pairs to the target.

Oligonucleotide 5 (SEQ ID NO:5) binds to the same 21 base target sequence on the target oligonucleotide that oligonucleotides 1 and 2 together bind. Oligonucleotide 6, a 22mer (SEQ ID NO:6) and oligonucleotide 7, a 23mer (SEQ ID NO:7) have 1 and 2 mismatches, respectively, in position that correspond to 1 and 2 base separation when oligonucleotides 1+3 and 1+4 bind to the target sequence together. Oligonucleotide 8 (SEQ ID NO:8) is a 13mer control oligonucleotide that binds to the same sequence as oligonucleotides 2 and 3 adjacent to oligonucleotide 1 without a base separation between them.

To further improve the cooperative interactions of the oligonucleotides binding to the target sequence at abutting sites, oligonucleotides 1 and 2 were both extended at the site of junction with complementary sequences so that they form a duplex stem upon interaction with the target, as shown in FIG. 1B. This extended antisense dimerization domain is designed not to have any complementarity with the adjacent bases of the antisense oligonucleotide binding site on the target. Oligonucleotides 9-17 (SEQ ID NOS:9-17) have an extended sequence on either the 5'- or 3'-end of the

-19-

binding sequence, which forms a duplex stem between the two oligonucleotides when they bind to adjacent sites on the target (FIG. 1B). This extended antisense dimerization domain has no complementarity with the target sequence. Oligonucleotides 9 and 14 form a 3 base pair stem. Oligonucleotides 10 and 14 have the same length of extended antisense dimerization domain but with one base separating the two target sites of the binding oligonucleotide pair. Oligonucleotide pairs 11+15, 12+16, and 13+17 bind to the same length of the sequence on the target as oligonucleotide pair 10+14 but with 4, 5, and 7 base pair extended antisense dimerization domains, respectively.

The initial evidence for cooperative binding of oligonucleotides 1 and 2 to their target sequence comes from thermal melting studies. TABLE 2 shows thermal melting data of the duplexes of these oligonucleotides individually and together with other corresponding oligonucleotides (FIG. 2). When oligonucleotides 1 and 2 bound side by side to the target, the resulting duplex has a T_m of 47.8°C. Duplexes of oligonucleotides 1+3 and 1+4 with the target sequence have T_m s of 44.4°C and 46°C, respectively. The oligonucleotides 1 and 3 bind to the target with a 1 base gap between them, and the oligonucleotides 1 and 4 bind to the target with a 2 base gap between them. The T_m of the duplex formed by oligonucleotides 1 and 2 together with the target is more than the average of the duplexes formed by 1 and 2 individually with the target sequence (TABLE 2).

35

-20-

TABLE 2

Oligos SEQ ID (NO:	Complex ^{a,b}	T _m , °C
1	CTAGAAGGAGAGAGATGGGTGCGAGAG CCCACGCTC	49.1
2	CTAGAAGGAGAGAGATGGGTGCGAGAG TTCCTCTCTCTA	43.4
3	CTAGAAGGAGAGAGATGGGTGCGAGAG CTTCCTCTCTCT	43.6
4	CTAGAAGGAGAGAGATGGGTGCGAGAG TCTTCCTCTCTC	45.0
5	CTAGAAGGAGAGAGATGGGTGCGAGAG TTCCTCTCTCTACCCACGCTC	67.7
6	CTAGAAGGAGAGAGATGGGTGCGAGAG CTTCCTCTCTCTG <u>CCC</u> ACGCTC	64.2
7	CTAGAAGGAGAGAGATGGGTGCGAGAG TCTTCCTCTCTC <u>CG</u> CCCACGCTC	59.9
1+2	CTAGAAGGAGAGAGATGGGTGCGAGAG TTCCTCTCTCTACCCACGCTC	47.8
1+3	CTAGAAGGAGAGAGATGGGTGCGAGAG CTTCCTCTCTCT CCCACGCTC	44.4
1+4	CTAGAAGGAGAGAGATGGGTGCGAGAG TCTTCCTCTCTC CCCACGCTC	45.9
1+8	CTAGAAGGAGAGAGATGGGTGCGAGAG CTTCCTCTCTCTACCCACGCTC	50.5

^a = underlined bases represent mismatches

^b = Target (SEQ ID NO:21) is bolded and is 5' → 3';
cooperative oligonucleotides are 3' → 5'.

-21-

In contrast, in the latter two cases (1+3 and 1+4), the T_m s are below the average of the two individual oligonucleotides in experiment. Further, in the case of the duplex formed with oligonucleotides 1+2 a sharp, single, cooperative transition was noticed (FIG. 2B). However, in the cases of the duplexes formed with 1+3 and 1+4, melting transitions were broad (FIG. 2B). This indicates that the two short oligonucleotides 1 and 2 targeted to two adjacent sites bind in a cooperative fashion, whereas those which bind leaving a one or two base gap between them do not interact cooperatively.

The duplex of oligonucleotide 5 which binds to the entire 21 base length has a T_m of 67.7°C. The duplex of oligonucleotide 6 (SEQ ID NO:6), a 22-mer with a mismatch in place that corresponds to one base gap between oligonucleotides 1 and 3, has a T_m of 64.2°C. Similarly, the duplex of oligonucleotide 7 (SEQ ID NO:7), a 23mer with two mismatches in a position that corresponds to the two base gap between oligonucleotides 1 and 4, has a T_m of 59.9°C. The lower melting temperatures of oligonucleotides 6 and 7 which bind to the target with one or two base mismatches indicate that these oligonucleotides can bind to a number of sites other than the perfectly matched target site at physiological temperatures. Thus, sequence specificity is decreasing.

Thermal melting studies of the duplexes of the oligonucleotides 9-17 demonstrates that the binding of these tandem oligonucleotides is further facilitated by the duplex stem (i.e., antisense dimerization domain) formed by extending the antisense dimerization domain. The stability of the ternary complex formed increases with an increase in the number of base pairs in the antisense dimerization domain, as shown in TABLE 3.

-22-

TABLE 3

Oligos (SEQ ID NOS:) ^a	Complex ^b	T _m , °C
10+14	CTAGAAGGAGAGAGATGGGTGCGAGAG CTCCTCTCTCT CCCACGCTC G C G C C G	45.9
11+15	CTAGAAGGAGAGAGATGGGTGCGAGAG CTCCTCTCTCT CCCACGCTC G C G C C G C G	47.3
12+16	CTAGAAGGAGAGAGATGGGTGCGAGAG CTCCTCTCTCT CCCACGCTC G C G C C G C G G C	48.4
13+1	CTAGAAGGAGAGAGATGGGTGCGAGAG CTCCTCTCTCT CCCACGCTC G C G C C G C G G C C G G C	53.2
9+14	CTAGAAGGAGAGAGATGGGTGCGAGAG TTCCTCTCTCTACCCACGCTC GC GC CG	47.9

^a 3' cooperative oligonucleotide then 5' cooperative oligonucleotide.

^b Target (SEQ ID NO:21) is bolded and is 5' → 3'; complementary cooperative oligonucleotides are 3' → 5'

For example, the double helical complexes with 3 base pair (oligonucleotides 10+14), 4 base pair

-23-

(oligonucleotides 11+15), 5 base pair
(oligonucleotides 12+16), and 7 base pair
(oligonucleotides 13+17) antisense dimerization
domains gave Tms of 45.9°C, 47.3°C, 48.4°C and
5 53.2°C, respectively. Further increases in duplex
stem length results in the formation of a stable
complex between the two tandem oligonucleotides in
the absence of the target sequence, an occurrence
which is not desirable. In all the cases, a sharp
10 cooperative single melting transition was observed
(FIG. 3).

Modified cooperative oligonucleotides were
studied for their antisense abilities. For example,
15 phosphorothioate internucleotide-linked forms of
cooperative oligonucleotides were studied for their
ability to activate RNase H. RNase H is an enzyme
that recognizes RNA-DNA heteroduplexes and
hydrolyses the RNA component of the heteroduplex
20 (Cedergren et al. (1987) *Biochem. Cell Biol.* 65:677).
Some studies have shown that antisense
oligonucleotides have less transition inhibition
activity in RNase H-free systems than in systems
where RNase H is present (Haeuptle et al. (1986)
25 *Nucleic Acids Res.* 14:1427-14448; Minshull et al. (1986)
Nucleic Acids Res. 14:6433-6451), or when the chemical
modification on antisense oligonucleotide is unable
to evoke RNase H activity (Maher III et al. (1988)
Nucl. Acids Res. 16:3341-3358; Leonetti et al. (1988)
30 *Gene* 72:323-332). In addition, it has also been
showed that a 4 to 6 base pair long hybrid is
sufficient to evoke RNase H activity.

-24-

A 39mer RNA target sequence (SEQ ID NO:22) which encodes a portion of the HIV-1 *gag* gene (TABLE 1) was synthesized to study the RNase H activation property of modified cooperative oligonucleotides of the invention. As per the design, modified oligonucleotides 1, 10, and 17 bind to a 9 base site on the 3'-side of the binding site of the target, and modified oligonucleotides 2, 13, and 14 bind on the 5'-side of the target adjacent to the binding site of the former oligonucleotide. Oligonucleotide 5 binds to the entire length of the 21 bases on the target. Oligonucleotides 6, 7, 18 and 19 contained mismatches.

15 An autoradiogram showing the RNase H hydrolysis pattern of the RNA target in the absence and presence of oligonucleotides of the invention is shown in FIGS. 4A and 4B. As expected, in experiments 2 and 5 (FIG. 4A), and in experiment 2 (FIG. 4B), hydrolytic activity is observed towards the 3'-end of the target RNA (lower half of the autoradiogram) in which oligonucleotides 1, 14, and 17, respectively, are present. Similarly, in experiments 3 and 6 (FIG. 4A) and in experiment 3 (FIG. 4B), RNA degradation bands are present only in the upper half of the autoradiogram, indicating the binding of oligonucleotides 2, 10, and 13, respectively, on the 5'-side of the target. When combinations of oligonucleotides are present (i.e., 1+2, 10+14, and 13+17) in experiments 4 and 7 (FIG. 4A) and in experiment 4 (FIG. 4B), the RNase H degradation pattern obtained is very similar to the one observed in the case of control oligonucleotide 5 in experiment 1 (FIGS. 5A and 5B). This clearly indicates that the new short tandem cooperative

-25-

oligonucleotides of the invention bind to the target RNA as expected with sequence specificity and evoke RNase H activity.

5 To further understand sequence specificity of
the cooperative oligonucleotides versus longer
oligonucleotides, two short oligonucleotides
analogous to oligonucleotide 1 having one and two
mismatches, oligonucleotides 18 (SEQ ID NO:18) and
10 19 (SEQ ID NO:19), were synthesized and studied for
RNase H activation in comparison to oligonucleotides
23 and 24. FIG. 5 shows the RNase H hydrolytic
pattern of target RNA in the presence of the
mismatched oligonucleotides. Oligonucleotide 23
15 (SEQ ID NO:23) with 1 mismatch (experiment 2) shows
the same RNase H degradation pattern as completely
matched oligonucleotide 5 (experiment 1).
Oligonucleotide 24 (SEQ ID NO:24) with two
mismatches (experiment 3) shows little or no RNA
20 hydrolysis in the middle of the binding site, where
the mismatches are located. However, on either side
of the mismatches the degradation pattern is exactly
like that found with oligonucleotide 5 which has no
mismatches. This clearly indicates that, in spite
25 of the two mismatches, oligonucleotide 24 binds to
the target strongly enough to activate RNase H.
Oligonucleotide 18 with one mismatch (experiment 5)
shows little or no RNA degradation compared to
oligonucleotide 1 (experiment 4). However, it
30 appears that oligonucleotide 18 has a strong binding
site on the 5'-end of the RNA target as indicated by
the RNA degradation bands towards the 5'-end of the
RNA. No digestion of the 3'-end of the RNA target
and little digestion of the 5'-end was observed with
35 oligonucleotide 19, which has two mismatches

-26-

(experiment 6). This clearly demonstrates that the new cooperative oligonucleotides bind with sequence specifically.

5 Representative modified cooperative
oligonucleotides of the invention were also studied
for their HIV-1 virus inhibition properties in cell
cultures. The results using phosphorothioate
cooperative oligonucleotides are shown in FIG. 6 as
10 a graph of percent virus inhibition versus
concentration of the oligonucleotide(s) and FIG. 7.
Oligonucleotide 5, a 21mer that is 4 bases shorter
than oligonucleotide 20, demonstrated little or no
significant activity up to a 15 μ M concentration.
15 Similarly, the combination of oligonucleotides 1+2,
which bind to the same sequence on the target as
oligonucleotide 5, also failed to show much
activity. The IC_{50} for oligonucleotide 20 in the
same assay system was about 0.55 μ M. In contrast, a
20 pronounced synergistic effect is observed with
oligonucleotide combination 13+17 which forms a 7
base pair dimerization duplex stem. This
oligonucleotide combination showed activity close to
oligonucleotide 20, with an IC_{50} value of about 4.0
25 μ M. The combination 10+4, which forms a three base
pair extended dimerization stem, showed about 15%
virus inhibition at 4 μ M concentration
(FIG. 7). Combination 12+16, with a five base
extended dimerization domain, showed about 25% viral
30 inhibition at the same concentration (FIG. 7).
Thus, the inhibition of HIV-1 virus progression by
combinations of oligonucleotides is higher than the
average of either oligonucleotide of the pair tested
alone. Note that the concentration of each
35 oligonucleotide in a combination is half that of the

-27-

individual oligonucleotide tested alone. For example, the concentration of oligonucleotides 13 and 17 is 2 plus 2, to a total concentration of 4 μ M, whereas the concentration of oligonucleotide 17, when it was tested alone, was 4 μ M. The other oligonucleotides studied individually or in combinations did not show significant activity even up to 10 μ M concentration (FIG. 7). The oligonucleotides 9+14, which form a 3 base pair duplex stem without a base separation between the binding oligonucleotides on the target, showed comparable activity to that of the combination of oligonucleotides 12 and 16, which form a 5 base pair duplex stem but with a one base separation. This result correlates well with the T_m data (Table 3).

The oligonucleotide combinations with an extended dimerization domain inhibited HIV much more efficiently than oligonucleotide 5 or the combination of oligonucleotides 1 and 2. FIG. 8 shows the relationship between HIV-1 inhibition and T_m of the complex formed. The oligonucleotide combination 13 and 17, which forms a 7 base pair antisense duplex stem, showed significantly greater activity relative to the other combinations of oligonucleotides, which form 3, 4, and 5 base pair duplex stems and oligonucleotide 5, a 21-mer.

These results demonstrate that modified cooperative oligonucleotides with dimerization domains have an enhanced ability to inhibit the expression of the target gene.

As described above, cooperative oligonucleotides of the invention can be directed to

-28-

any target oligonucleotide. Other non-limiting examples of such cooperative oligonucleotides include those specific for influenza nucleic acid targets which are shown below in Table 4.

5

TABLE 4

10	Target Oligo (SEQ ID NO:)	Complex ^{a,b}	Cooperative Oligo (SEQ ID NO:)
	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTTACCT	41, 40
15	39	TCCATTCAAATGGTTTGCCTGC AGGTAAGTTTACCAAACGGACG	42, 43
	39	TCCATTCAAATGGTTTGCCTGC AGGTAAGTTTACCAAACGGACG	44, 45
20		GC CG TA CG GC	
25	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTTACCT	46, 47
30		GC TA CG GC CG	
35	39	TCCATTCAAATGGTTTGCCTGC AGGTAAGTTTACCAAACGGACG	48
	39	TCCATTCAAATGGTTTGCCTGC AGGTAAGTTTACCAAACGGACG	49
40	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTTACCT	50
	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTTACCT	51
45	39	TCCATTCAAATGGTTTGCCTGC AGGTAAGTTTACCAAACGGACG	44, 52
50		GC CG TA CG GC	
55	39	TCCATTCAAATGGTTTGCCTGC AGGTAAGTTTACCAAACGGACG	53, 54
60		GC CG TA CG GC	

-29-

TABLE 4 (con't)

5	Target Oligo (SEQ ID NO:)	Complex ^{a,b}	Cooperative Oligo (SEQ ID NO:)
10	39	TCCATTCAAATGGTTTGCCTGC AGGTATGTTTACCAAAAGGACG GC CG TA CG GC	53, 52
15	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACATACCT GC TA CG GC CG	46, 55
20	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTACCT GC TA CG GC CG	56, 47
25	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTACCT GC TA CG GC CG	56, 55
30	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTACCT GC TA CG GC CG	
35	38	GCAGGCAAACCATTTGAATGGA CGTCCGTTTGGTAAACTACCT GC TA CG GC CG	
40	^a = underlined bases represent mismatches ^b = Target is bolded and is 5' → 3'; cooperative oligonucleotides are 3' → 5'.		

45

In addition, sequence-specific and cooperative binding of short oligonucleotides that bind to adjacent sites are useful to target sequences with point mutations specifically. In addition, undesirable non-sequence specific effects can be reduced by using at least two short oligonucleotides that can bind to a longer target sequence rather than one long oligonucleotide that binds to the same length of the target sequence. For example, long oligonucleotides that contain a modified backbone, such as phosphorothioates, activate complement, which have adverse cardiovascular effects (Galbraith

-30-

et al. (1994) *Antisense Res. Dev.* 4:201-207; and Cornish
et al. (1993) *Pharmacol. Commun.* 3:239-247). In
conclusion, combination oligonucleotides represent
an alternative therapeutic strategy to the use of a
5 single oligonucleotide, in cases in which use of the
latter is limited by concentration and chain length
constraints, and the associated problems of toxicity
and production costs.

10 The synthetic cooperative oligonucleotides of
the invention also may be used to identify the
presence of the nucleic acids of a particular virion
or bacteria in cell cultures, for example, by
labelling the oligonucleotide and screening for
15 double-stranded, labelled DNA in the cells by *in situ*
hybridization or some other art-recognized detection
method.

In addition, the function of various genes in
20 an animal, including those essential to animal
development can be examined using the cooperative
oligonucleotides of the invention. Presently, gene
function can only be examined by the arduous task of
making a "knock out" animal such as a mouse. This
25 task is difficult, time-consuming and cannot be
accomplished for genes essential to animal
development since the "knock out" would produce a
lethal phenotype. The present invention overcomes
the shortcomings of this model.

30

It is known that antisense oligonucleotides can
bind to a target single-stranded nucleic acid
molecule according to the Watson-Crick or the
Hoogsteen rule of base pairing, and in doing so,
35 disrupt the function of the target by one of several

-31-

mechanisms: by preventing the binding of factors required for normal transcription, splicing, or translation; by triggering the enzymatic destruction of mRNA by RNase H if a contiguous region of

5 deoxyribonucleotides exists in the oligonucleotide, and/or by destroying the target via reactive groups attached directly to the antisense oligonucleotide.

Thus, because of the properties described

10 above, such oligonucleotides are useful therapeutically by their ability to control or down-regulate the expression of a particular gene in a cell, e.g., in a cell culture or in an animal, according to the method of the present invention.

15 The cooperative oligonucleotides of the invention may also be used to inhibit transcription of any gene in a cell, including a foreign gene. For example, the cooperative oligonucleotides as

20 provided by the invention may be used to inhibit the expression of HIV genes within infected host cells and thus to inhibit production of HIV virions by those cells. The synthetic oligonucleotides of the invention are thus useful for treatment of HIV

25 infection and AIDS in mammals, particularly the treatment of mammals used as animal models to study HIV infection and AIDS. The synthetic oligonucleotides of the invention are also useful for treatment of humans infected with HIV and those

30 suffering from AIDS.

As discussed above, the synthetic oligonucleotides of the invention may be used as a pharmaceutical composition when combined with a

35 pharmaceutically acceptable carrier. The term

-32-

"pharmaceutically acceptable" means a non-toxic material that does not interfere with the effectiveness of the biological activity of the active ingredient(s). The characteristics of the carrier will depend on the route of administration. Such a composition may contain, in addition to the synthetic oligonucleotide and carrier, diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. The pharmaceutical composition of the invention may also contain other active factors and/or agents which enhance inhibition of virus or bacterial production by infected cells. For example, combinations of synthetic oligonucleotides, each of which inhibits transcription of a different HIV gene, may be used in the pharmaceutical compositions of the invention. The pharmaceutical composition of the invention may further contain nucleotide analogs such as azidothymidine, dideoxycytidine, dideoxyinosine, and the like. Such additional factors and/or agents may be included in the pharmaceutical composition to produce a synergistic effect with the synthetic oligonucleotide of the invention, or to minimize side-effects caused by the synthetic oligonucleotide of the invention. Conversely, the synthetic oligonucleotide of the invention may be included in formulations of a particular anti-HIV factor and/or agent to minimize side effects of the anti-HIV factor and/or agent.

30

The pharmaceutical composition of the invention may be in the form of a liposome in which the synthetic oligonucleotides of the invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as

35

-33-

lipids which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers which are in aqueous solution. Suitable lipids for liposomal formulation include, without
5 limitation, monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of such liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S.
10 Patent No. 4,235,871; U.S. Patent No. 4,501,728; U.S. Patent No. 4,837,028; and U.S. Patent No. 4,737,323.

The pharmaceutical composition of the invention
15 may further include compounds which enhance delivery of oligonucleotides into cells, as described in WO 95/32739.

As used herein, the term "therapeutically
20 effective amount" means the total amount of each active component of the pharmaceutical composition or method that is sufficient to show a meaningful patient benefit, e.g., healing of chronic conditions characterized by HIV and associated infections and
25 complications or by other viral infections or increase in rate of healing of such conditions. When applied to an individual active ingredient, administered alone, the term refers to that ingredient alone. When applied to a combination,
30 the term refers to combined amounts of the active ingredients that result in the therapeutic effect, whether administered in combination, serially or simultaneously.

-34-

In practicing the method of treatment or use of the present invention, a therapeutically effective amount of one or more of the synthetic oligonucleotide of the invention is administered to a mammal infected with HIV. The synthetic oligonucleotide of the invention may be administered in accordance with the method of the invention either alone or in combination with other therapies such as treatments employing cytokines, lymphokines, other hematopoietic factors, other anti-viral agents, and the like. When co-administered with one or more cytokines, lymphokines or other hematopoietic factors, other anti-viral agents, the synthetic oligonucleotide of the invention may be administered either simultaneously with the cytokine(s), lymphokine(s), other hematopoietic factor(s), other antiviral agents, and the like, or sequentially. If administered sequentially, the attending physician will decide on the appropriate sequence of administering the synthetic oligonucleotide of the invention in combination with cytokine(s), lymphokine(s), other hematopoietic factor(s), anti-viral agents, and the like.

Administration of the synthetic oligonucleotide of the invention used in the pharmaceutical composition or to practice the method of the present invention can be carried out in a variety of conventional ways, such as oral ingestion, inhalation, or cutaneous, subcutaneous, or intravenous injection. Intravenous administration to the patient is preferred.

When a therapeutically effective amount of synthetic oligonucleotide of the invention is

-35-

administered orally, the synthetic oligonucleotide will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the
5 invention may additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5 to 95% synthetic oligonucleotide and preferably from about 25 to 90% synthetic oligonucleotide. When
10 administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, sesame oil, or synthetic oils may be added. The liquid form of the pharmaceutical composition may
15 further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from
20 about 0.5 to 90% by weight of the synthetic oligonucleotide and preferably from about 1 to 50% synthetic oligonucleotide.

When a therapeutically effective amount of
25 synthetic oligonucleotide of the invention is administered by intravenous, cutaneous or subcutaneous injection, the synthetic oligonucleotide will preferably be in the form of a pyrogen-free, parenterally acceptable aqueous
30 solution. The preparation of such parenterally acceptable solutions, having due regard to pH, isotonicity, stability, and the like, is within the skill in the art. A preferred pharmaceutical composition for intravenous, cutaneous, or
35 subcutaneous injection should contain, in addition

-36-

to the synthetic oligonucleotide, an isotonic vehicle such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or
5 other vehicle as known in the art. The pharmaceutical composition of the present invention may also contain stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art.

10

The amount of synthetic oligonucleotide in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated, and on the nature of prior
15 treatments which the patient has undergone. Ultimately, the attending physician will decide the amount of synthetic oligonucleotide with which to treat each individual patient. Initially, the attending physician will administer low doses of the
20 synthetic oligonucleotide and observe the patient's response. Larger doses of synthetic oligonucleotide may be administered until the optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is
25 contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 1 ng to about 100 mg of synthetic oligonucleotide per kg body weight.

30

The duration of intravenous therapy using the pharmaceutical composition of the present invention will vary, depending on the severity of the disease being treated and the condition and potential
35 idiosyncratic response of each individual patient.

-37-

It is contemplated that the duration of each application of the synthetic oligonucleotide will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately, the attending physician will decide on the appropriate duration of intravenous therapy using the pharmaceutical composition of the present invention.

The following examples illustrate the preferred modes of making and practicing the present invention, but are not meant to limit the scope of the invention since alternative methods may be utilized to obtain similar results.

EXAMPLES

1. Cooperative Oligonucleotide Synthesis

Cooperative oligodeoxyribonucleotides were synthesized on a Milligen 8700 DNA synthesizer using β -cyanoethylphosphoramidite chemistry (*Meth. Mol. Biol.* (1993) Vol. 20 (Agrawal (ed.) Humana Press, Totowa, NJ, pp. 33-61) on a (500 Å controlled pore glass solid support). Monomer synthons and other DNA synthesis reagents were obtained from Milligen Biosearch (Burlington, MA). After the synthesis and deprotection, oligonucleotides were purified on reverse phase (C_{18}) HPLC, detritylated, desalted (Waters C_{18} sep-pack cartridges (Waters, Milford, MA), and checked for purity by polyacrylamide gel electrophoresis (Manniatitis et al. in *Molecular Cloning (A Laboratory Manual)*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY). Cooperative oligoribonucleotides and hybrids (RNA/DNA) cooperative oligonucleotides are prepared according

-38-

to the method(s) of Meteleev et al. (*FEBS. Lett.* (1988) 226:232-234; and Atabekov et al. (1988) *FEBS. Lett.* 232:96-98.

5 Cooperative phosphorothioate oligonucleotides for RNase H and tissue culture experiments were synthesized as above but using sulfurizing agent as oxidant instead of normal iodine oxidant. Post-synthetic processing was carried out exactly as
10 above but desalting was performed by dialysis for 72 hours against double distilled water.

Other modified forms of the cooperative oligonucleotides are prepared as described in
15 Agrawal (ed.) (*Meth. Mol. Biol.*, Vol. 20, *Protocols for Oligonucleotides and Analogs*, (1993) Humana Press, Totowa, NJ).

2. UV Melting Studies

20

UV melting experiments were carried out in 150 mM sodium chloride, 10 mM sodium dihydrogen phosphate, and 2 mM magnesium chloride, pH 7.4 buffer. The oligonucleotide concentration was 0.36
25 μ M as single strand. The oligonucleotides were mixed in buffer, heated to 95°C, cooled down to room temperature, and left at 4°C overnight. Thermal denaturation profiles were recorded at 260 nm at a heating rate of 0.5°C/min on a spectrophotometer
30 (Perkin-Elmer Lambda2, (Norwalk CT) equipped with a peltier thermal controller and attached to a personal computer for data collection. The (T_m) melting temperatures were measured from first derivative plots (dA/dT vs T). Each value is an

-39-

average of two separate runs and the values are within $\pm 1.0^{\circ}\text{C}$ range.

3. RNase H Assay

5

An RNA target (SEQ ID NO:22) was labelled at its 3'-end using terminal transferase and [α - ^{32}P]ddATP (Amersham, (Arlington Heights, IL) using standard protocols (Manniatis et al. in *Molecular Cloning, A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY). End-labelled RNA (3000-5000 cpm) was incubated with 1 to 1.5 ratio of the oligonucleotides in 30 μl of 20 mM Tris-HCl, pH 7.5, 10 mM MgCl_2 , 10 mM KCl, 0.1 mM DTT, 5% sucrose (w/v), and 40 units of RNasin (Promega, Madison, WI) at 4°C overnight. An aliquot (7 μl) was taken out as control, 1 μl (0.8 unit) of *E. coli* RNase H (Promega, Madison, WI) was added to the remaining reaction mixture and incubated at room temperature. Aliquots (7 μl) were taken out at different time intervals. The samples were then analyzed on a 7 M urea 20% polyacrylamide gel. After the electrophoresis, an autoradiogram was developed by exposing the gel to Kodak X-Omat AR film at -70°C .

4. Antiviral Assay

The effect of the antisense oligonucleotides on the replication of HIV-1 during an acute infection was determined. The test system is a modification of the standard cytopathic effect (CPE)-based MT-2 cell assay (Posner et al. (1991) *J. Immunol.* 146:4325; Pawels et al. (1988) *J. Virol. Methods* 20:309; Mosmann

-40-

(1983) *J. Immunol. Methods* 65:55). Briefly, serial dilutions of antisense oligonucleotides synthesized as described above, or the combinations of such oligonucleotides, were prepared in 50 μ M L-glutamine, 100 U/ml penicillin, 100 μ g/ml streptomycin), in triplicate, in 96-well plates. Virus, (HIV-1 IIIB originally obtained from Dr. Robert Gallo, NCI (Popovic et al. (1984) *Science* 224:497) and propagated in H9 cells (Gazdar et al. (1980) *Blood* 55:409) by the method of Vujcic (*J. Infect. Dis.* (1988) 157:1047), diluted to contain a 90% cytopathic effect (CPE) dose of virus in 50 μ l, was added followed by 100 μ l of 4×10^5 /ml MT-2 cells (Harada et al. (1985) *Science* 229:563) in complete medium. The plates were incubated at 37°C in 5% CO₂, for 5 days. 3-[4, 5-dimethylthiazol-2-yl]-2, 5-diphenyltetrazolium bromide; thiazoyl blue (MTT) dye (Sigma, St. Louis, MO) was added and quantitated at OD₅₄₀-OD₆₉₀ as described (Posner et al. (1991) *J. Immunol.* 146:4325). Percent viral inhibition was calculated by the formula: (experimental-virus control)/(medium control-virus control) x 100.

EQUIVALENTS

25

Those skilled in the art will recognize, or be able to ascertain, using no more than routine experimentation, numerous equivalents to the specific substances and procedures described herein. Such equivalents are considered to be within the scope of this invention, and are covered by the following claims.

-41-

SEQUENCE LISTING

(1) GENERAL INFORMATION:

- (i) APPLICANT: Kandimalla, Ekambar R.
Agrawal, Sudhir
- (ii) TITLE OF INVENTION: COOPERATIVE OLIGONUCLEOTIDES
- (iii) NUMBER OF SEQUENCES: 57
- (iv) CORRESPONDENCE ADDRESS:
 - (A) ADDRESSEE: Lappin & Kusmer
 - (B) STREET: 200 State Street
 - (C) CITY: Boston
 - (D) STATE: MA
 - (E) COUNTRY: USA
 - (F) ZIP: 02109
- (v) COMPUTER READABLE FORM:
 - (A) MEDIUM TYPE: Floppy disk
 - (B) COMPUTER: IBM PC compatible
 - (C) OPERATING SYSTEM: PC-DOS/MS-DOS
 - (D) SOFTWARE: PatentIn Release #1.0, Version #1.30
- (vi) CURRENT APPLICATION DATA:
 - (A) APPLICATION NUMBER: US
 - (B) FILING DATE:
 - (C) CLASSIFICATION:
- (viii) ATTORNEY/AGENT INFORMATION:
 - (A) NAME: Kerner, Ann-Louise
 - (B) REGISTRATION NUMBER: 33,523
 - (C) REFERENCE/DOCKET NUMBER: HYZ-027PCT
- (ix) TELECOMMUNICATION INFORMATION:
 - (A) TELEPHONE: 617-330-1300
 - (B) TELEFAX: 617-330-1311

(2) INFORMATION FOR SEQ ID NO:1:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 9 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear
- (ii) MOLECULE TYPE: cDNA
- (iii) HYPOTHETICAL: NO

-42-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:1:

CTCGCACCC

9

(2) INFORMATION FOR SEQ ID NO:2:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:2:

ATCTCTCTCC TT

12

(2) INFORMATION FOR SEQ ID NO:3:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:3:

TCTCTCTCCT TC

12

(2) INFORMATION FOR SEQ ID NO:4:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 12 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-43-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:4:

CTCTCTCCTT CT

12

(2) INFORMATION FOR SEQ ID NO:5:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:5:

CTCGCACCCA TCTCTCTCCT T

21

(2) INFORMATION FOR SEQ ID NO:6:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 22 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:6:

CTCGCACCCG TCTCTCTCCT TC

22

(2) INFORMATION FOR SEQ ID NO:7:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 23 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-44-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:7:

CTCGCACCCG CCTCTCTCCT TCT

23

(2) INFORMATION FOR SEQ ID NO:8:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:8:

ATCTCTCTCC TTC

13

(2) INFORMATION FOR SEQ ID NO:9:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:9:

CGGATCTCTC TCCTT

15

(2) INFORMATION FOR SEQ ID NO:10:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 15 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-45-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:10:

CGGTCTCTCT CCTTC

15

(2) INFORMATION FOR SEQ ID NO:11:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:11:

CCGGTCTCTC TCCTTC

16

(2) INFORMATION FOR SEQ ID NO:12:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:12:

GCCGGTCTCT CTCCTTC

17

(2) INFORMATION FOR SEQ ID NO:13:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 19 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-46-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:13:

GCGCCGGTCT CTCTCCTTC

19

(2) INFORMATION FOR SEQ ID NO:14:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 12 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:14:

CTCGCACCCC CG

12

(2) INFORMATION FOR SEQ ID NO:15:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 13 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:15:

CTCGCACCCC CGG

13

(2) INFORMATION FOR SEQ ID NO:16:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 14 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-47-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:16:

CTCGCACCCC CGGC

14

(2) INFORMATION FOR SEQ ID NO:17:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 16 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:17:

CTCGCACCCC CGGCGC

16

(2) INFORMATION FOR SEQ ID NO:18:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:18:

CTCTCACCC

9

(2) INFORMATION FOR SEQ ID NO:19:

- (i) SEQUENCE CHARACTERISTICS:
 (A) LENGTH: 9 base pairs
 (B) TYPE: nucleic acid
 (C) STRANDEDNESS: single
 (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

-48-

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:19:

CTCTCAACC

9

(2) INFORMATION FOR SEQ ID NO:20:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 25 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:20:

CTCTCGCACC CATCTCTCTC CTTCT

25

(2) INFORMATION FOR SEQ ID NO:21:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 27 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: DNA (genomic)

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:21:

CTAGAAGGAG AGAGATGGGT GCGAGAG

27

(2) INFORMATION FOR SEQ ID NO:22:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 39 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(iii) HYPOTHETICAL: NO

-49-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:22:

AGAAGGAGAG AGAUGGGUGC GAGAGCGUCA GUAUUAAGC

39

(2) INFORMATION FOR SEQ ID NO:23:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:23:

CTCTCACCCA TCTCTCTCCT T

21

(2) INFORMATION FOR SEQ ID NO:24:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 21 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: YES

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:24:

CTCTCAACCA TCTCTCTCCT T

21

(2) INFORMATION FOR SEQ ID NO:25:

(i) SEQUENCE CHARACTERISTICS:

- (A) LENGTH: 35 base pairs
- (B) TYPE: nucleic acid
- (C) STRANDEDNESS: single
- (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: RNA

(iii) HYPOTHETICAL: NO

-50-

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:25:

GGAGGCUAGA AGGAGAGAGA UGGGUGCGAG AGCGU

35

(2) INFORMATION FOR SEQ ID NO:26:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:26:

GACCACTTCT CTTTCT

16

(2) INFORMATION FOR SEQ ID NO:27:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 21 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:27:

GAATGAGCCA TCTCAGTGGT C

21

(2) INFORMATION FOR SEQ ID NO:28:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-51-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:28:

CTCTCGCACC TCACTC

16

(2) INFORMATION FOR SEQ ID NO:29:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:29:

CCACTTCTCC TTCT

14

(2) INFORMATION FOR SEQ ID NO:30:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:30:

ATGAGCCATC TCAGTGG

17

(2) INFORMATION FOR SEQ ID NO:31:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 14 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-52-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:31:

CTCTCGCACC TCAC

14

(2) INFORMATION FOR SEQ ID NO:32:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:32:

ACTTCTCCTT CT

12

(2) INFORMATION FOR SEQ ID NO:33:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 13 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:33:

GAGCCATCTC AGT

13

(2) INFORMATION FOR SEQ ID NO:34:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-53-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:34:

CTCTCGCACC TC

12

(2) INFORMATION FOR SEQ ID NO:35:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:35:

TCTCCTTCT

9

(2) INFORMATION FOR SEQ ID NO:36:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 7 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:36:

CCATCTC

7

(2) INFORMATION FOR SEQ ID NO:37:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 9 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-54-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:37:

CTCTCGCAC

9

(2) INFORMATION FOR SEQ ID NO:38:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:38:

GCAGGCAAAC CATTGAATG GA

22

(2) INFORMATION FOR SEQ ID NO:39:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: NO

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:39:

TCCATTCAAA TGGTTTGCCT GC

22

(2) INFORMATION FOR SEQ ID NO:40:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 11 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-55-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:40:

TCCATTCAAA T

11

(2) INFORMATION FOR SEQ ID NO:41:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:41:

GGTTTGCCTG C

11

(2) INFORMATION FOR SEQ ID NO:42:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:42:

ATTTGAATGG A

11

(2) INFORMATION FOR SEQ ID NO:43:

- (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 11 base pairs
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-56-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:43:

GCAGGCAAAC C

11

(2) INFORMATION FOR SEQ ID NO:44:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:44:

GCTCGATTTG AATGGA

16

(2) INFORMATION FOR SEQ ID NO:45:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:45:

GCAGGCAAAC CCGAGC

16

(2) INFORMATION FOR SEQ ID NO:46:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-57-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:46:

CGCTGGGTTT GCCTGC

16

(2) INFORMATION FOR SEQ ID NO:47:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:47:

TCCATTCAAA TCAGCG

16

(2) INFORMATION FOR SEQ ID NO:48:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:48:

GCAGGAAAAC CATTGAATG GA

22

(2) INFORMATION FOR SEQ ID NO:49:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 22 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

-58-

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:49:

GCAGGAAAAC CATTGTATG GA

22

(2) INFORMATION FOR SEQ ID NO:50:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:50:

TCCATACAAA TGGTTTGCCT GC

22

(2) INFORMATION FOR SEQ ID NO:51:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 22 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:51:

TCCATACAAA TGGTTTCCCT GC

22

(2) INFORMATION FOR SEQ ID NO:52:

- (i) SEQUENCE CHARACTERISTICS:
 - (A) LENGTH: 16 base pairs
 - (B) TYPE: nucleic acid
 - (C) STRANDEDNESS: single
 - (D) TOPOLOGY: linear

-59-

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:52:

GCAGGAAAAC CCGAGC

16

(2) INFORMATION FOR SEQ ID NO:53:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:53:

GCTCGATTTG TATGGA

16

(2) INFORMATION FOR SEQ ID NO:54:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:54:

GCAGGCAAAC CCGAGC

16

(2) INFORMATION FOR SEQ ID NO:55:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

-60-

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:55:

TCCATACAAA TCAGCG

16

(2) INFORMATION FOR SEQ ID NO:56:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 16 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:56:

CGCTGGGTTT CCCTGC

16

(2) INFORMATION FOR SEQ ID NO:57:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 12 base pairs

(B) TYPE: nucleic acid

(C) STRANDEDNESS: single

(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: cDNA

(iii) HYPOTHETICAL: NO

(iv) ANTI-SENSE: YES

(xi) SEQUENCE DESCRIPTION: SEQ ID NO:57:

TCTCTCTCCT TC

12

-61-

What is claimed is:

1. A composition comprising at least a first synthetic cooperative oligonucleotide and a second synthetic cooperative oligonucleotide, each oligonucleotide comprising a region complementary to one of tandem, non-overlapping regions of a target nucleic acid, and a dimerization domain at a terminus of each of the oligonucleotides,
 - the dimerization domains of the oligonucleotides being complementary to each other, and
 - the target nucleic acid being an mRNA, a single-stranded viral DNA, or a single-stranded viral RNA.
2. The composition of claim 1 wherein the oligonucleotides are complementary to tandem regions of the target nucleic acid that are separated by 0 to 3 bases.
3. The composition of claim 1 wherein each of the oligonucleotides are about 9 to 25 nucleotides in length.
4. The composition of claim 1 wherein the dimerization domain of the first cooperative oligonucleotide is located at its 3' terminal portion, and is complementary to the dimerization domain of the second oligonucleotide which is located at its 5' terminal portion.
5. The composition of claim 1 wherein the dimerization domain of a first cooperative oligonucleotide is located at its 3' terminal portion, and is complementary to the dimerization domain of a second oligonucleotide which is located at its 3' terminal portion.

-62-

6. The composition of claim 1 wherein the dimerization domain of a first cooperative oligonucleotide is located at its 5' terminal portion, and is complementary to the dimerization domain of a second oligonucleotide which is located at its 5' terminal portion.
7. The composition of claim 1 wherein the dimerization domain of each of the oligonucleotides are 3 to 7 nucleotides in length.
8. The composition of claim 1 wherein the first and second regions of the nucleic acid that are separated by 0 to 3 bases.
9. The composition of claim 1 wherein at least one of the oligonucleotides is modified.
10. The composition of claim 9 wherein at least one of the oligonucleotides contains at least one non-phosphodiester internucleotide linkage.
11. The composition of claim 9 wherein at least one of the oligonucleotides contains at least one phosphorothioate internucleotide linkage.
12. The composition of claim 1 wherein first and second oligonucleotides are hybridized to the target nucleic acid and the dimerization domain of the first oligonucleotide is hybridized to the dimerization domain of the second oligonucleotide.

-63-

13. A duplex structure comprising a first synthetic cooperative oligonucleotide and a second synthetic cooperative oligonucleotide, each oligonucleotide comprising a region complementary to one of tandem, non-overlapping regions of a target nucleic acid,
5 the target nucleic acid being an mRNA, a single-stranded viral RNA, or a single-stranded viral DNA, and the first oligonucleotide having a terminal dimerization domain complementary and hybridized to a
10 dimerization domain of the second oligonucleotide when the first and second oligonucleotides are hybridized to the target nucleic acid.
14. The structure of claim 13 wherein each of the
15 oligonucleotides are about 9 to 25 nucleotides in length.
15. The structure of claim 13 wherein the dimerization domains of the first and second oligonucleotides each
20 comprise about 3 to 7 nucleotides.
16. The structure of claim 13 wherein the first and second oligonucleotides are complementary to one of
25 tandem regions of the target nucleic acid that are separated by 0 to 3 bases.
17. The structure of claim 13 hybridized to the target nucleic acid.

-64-

18. A composition comprising at least a first synthetic cooperative oligonucleotide, a second synthetic cooperative oligonucleotide, and a third synthetic cooperative oligonucleotide, each oligonucleotide
5 comprising a region complementary to one of tandem, non-overlapping regions of a target nucleic acid, and a dimerization domain at one or both termini of each of the oligonucleotides,

the dimerization domain of the first and second
10 oligonucleotide is at a single terminus of the first and second cooperative oligonucleotides, and a third cooperative oligonucleotide has a dimerization domain at both of its termini,

the dimerization domain of the first cooperative
15 oligonucleotide being complementary to one dimerization domain of the third cooperative oligonucleotide, and the dimerization domain of the second cooperative oligonucleotide being complementary to the second dimerization domain of the third cooperative
20 oligonucleotide, and

the target nucleic acid being an mRNA, a single-stranded viral DNA, or a single-stranded viral RNA.

19. The composition of claim 18 wherein the
25 oligonucleotides are complementary to tandem regions of the target nucleic acid that are separated by 0 to 3 bases.

20. The composition of claim 18 wherein each of the
30 oligonucleotides are about 9 to 25 nucleotides in length.

21. The composition of claim 18 wherein the dimerization domain of each of the oligonucleotides are 3 to 7 nucleotides in length.

35

22. The composition of claim 18 wherein at least one of the oligonucleotides is modified.

-65-

23. The composition of claim 22 wherein at least one of the oligonucleotides contains at least one non-phosphodiester internucleotide linkage.

5 24. The composition of claim 23 wherein at least one of the oligonucleotides contains at least one phosphorothioate internucleotide linkage.

25. The composition of claim 18 wherein the dimerization
10 domain of the first cooperative oligonucleotide is located at its 3' terminus and is complementary to the dimerization domain at the 5' terminus of the third oligonucleotide, and the dimerization domain of the second cooperative oligonucleotide is located at its 5'
15 terminus and is complementary to the dimerization domain at the 3' terminus of the third oligonucleotide.

26. The composition of claim 18 wherein the dimerization domain of the first cooperative oligonucleotide is
20 located at its 5' terminus and is complementary and hybridized to the dimerization domain at the 3' terminus of the third oligonucleotide, and the dimerization domain of the second cooperative oligonucleotide is located at its 3' terminus and is complementary and hybridized to
25 the dimerization domain at the 5' terminus of the third cooperative oligonucleotide.

-66-

27. A ternary structure comprising a first synthetic cooperative oligonucleotide, a second synthetic cooperative oligonucleotide, and a third synthetic cooperative oligonucleotide, each oligonucleotide
5 comprising a region complementary to one of tandem, non-overlapping regions of a target nucleic acid, and each comprising a dimerization domain at one or both of their termini,
the dimerization domain of the first oligonucleotide
10 being complementary and hybridized to a first dimerization domain at one terminus of the third oligonucleotide and the dimerization domain of the second oligonucleotide being complementary and hybridized to a second dimerization domain at the other terminus of the
15 third oligonucleotide when the first, second, and third oligonucleotides are hybridized to the target nucleic acid, and
the target nucleic acid being an mRNA, a single-stranded viral RNA, or a single-stranded viral DNA.
20
28. The structure of claim 27 wherein each of the oligonucleotides are about 9 to 25 nucleotides in length.
29. The structure of claim 27 wherein the dimerization
25 domains of the first and second oligonucleotides each comprise about 3 to 7 nucleotides.
30. The structure of claim 27 wherein the first and second oligonucleotides are complementary to one of
30 tandem regions of the target nucleic acid that are separated by 0 to 3 bases.
31. The ternary structure of claim 27 hybridized to the target nucleic acid.
35

-67-

32. A pharmaceutical formulation comprising the composition of claim 1 and a physiologically acceptable carrier.
- 5 33. A pharmaceutical formulation comprising the duplex structure of claim 13 and a physiologically acceptable carrier.
- 10 34. A pharmaceutical formulation comprising the composition of claim 18 and a physiologically acceptable carrier.
- 15 35. A pharmaceutical formulation comprising the composition of claim 27 and a physiologically acceptable carrier.
- 20 36. A method of inhibiting the expression of a nucleic acid comprising the step of contacting the nucleic acid with the composition of claim 1.
37. A method of inhibiting the expression of a nucleic acid comprising the step of contacting the nucleic acid with the composition of claim 13.
- 25 38. A method of inhibiting the expression of a nucleic acid comprising the step of contacting the nucleic acid with the composition of claim 18.
- 30 39. A method of inhibiting the expression of a nucleic acid comprising the step of contacting the nucleic acid with the composition of claim 27.
40. The method of claim 36 wherein the nucleic acid is a viral nucleic acid.
- 35 41. The method of claim 40 wherein the nucleic acid is an HIV DNA or an HIV RNA.

-68-

42. The method of claim 40 wherein the nucleic acid is an influenza DNA or influenza RNA.
43. The method of claim 37 wherein the nucleic acid is a
5 viral nucleic acid.
44. The method of claim 43 wherein the nucleic acid is an HIV DNA or an HIV RNA.
- 10 45. The method of claim 43 wherein the nucleic acid is an influenza DNA or influenza RNA.
46. The method of claim 38 wherein the nucleic acid is a viral nucleic acid.
15
47. The method of claim 46 wherein the nucleic acid is an HIV DNA or an HIV RNA.
48. The method of claim 46 wherein the nucleic acid is
20 an influenza DNA or influenza RNA.
49. The method of claim 39 wherein the nucleic acid is a viral nucleic acid.
- 25 50. The method of claim 49 wherein the nucleic acid is an HIV DNA or an HIV RNA.
51. The method of claim 49 wherein the nucleic acid is a viral nucleic acid.
30
52. A method of treating viral infection comprising the step of administering to the infected subject the pharmaceutical composition of claim 32.
- 35 53. A method of treating viral infection comprising the step of administering to the infected subject the pharmaceutical composition of claim 33.

-69-

54. A method of treating viral infection comprising the step of administering to the infected subject the pharmaceutical composition of claim 34.

- 5 55. A method of treating viral infection comprising the step of administering to the infected subject the pharmaceutical composition of claim 35.




FIG. 1A







FIG. 1B


2/10

*FIG. 1C*

3/10

**FIG. 2A****FIG. 2B**

4/10

**FIG. 3**

5/10





FIG. 4A

6/10



7/10

OLIGO. #	5	20	21	1	18	19
EXPT. #	1	2	3	4	5	6
TIME, MIN.	0 1 5 15	0 1 5 15	0 1 5 15	0 1 5 15	0 1 5 15	0 1 5 15

*FIG. 5*

8/10




FIG. 6





FIG. 7

10/10

**FIG. 8**

INTERNATIONAL SEARCH REPORT

International Application No

PC1/US 96/04605

A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/11 A61K31/70 C07H21/04

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO,A,95 01985 (LYNX THERAPEUTICS INC ;GRYAZNOV SERGEI M (US)) 19 January 1995 see page 8, line 28 - page 16, line 14 ---	1-51
A	GB,A,2 225 112 (ICI PLC) 23 May 1990 see claims 1-13 ---	1-14,32, 36
A	EP,A,0 185 494 (APPLIED BIOSYSTEMS) 25 June 1986 see claims 1-17 ---	1,13,32, 36
A	WO,A,91 06626 (GILEAD SCIENCES INC) 16 May 1991 see claims 1-12 ---	1,13,32, 36

	-/--	

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

- *T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- *X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- *Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- *&* document member of the same patent family

Date of the actual completion of the international search

4 September 1996

Date of mailing of the international search report

12.09.96

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax (+31-70) 340-3016

Authorized officer

Gurdjian, D

INTERNATIONAL SEARCH REPORT

International Application No

PC1/US 96/04605

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	WO,A,94 17086 (APOLLON INC ;YOON KYONGGEUN (US); LU MEIQING (US)) 4 August 1994 see figures 9,10 ---	1,13,32, 36
A	WO,A,94 23028 (HYBRIDON INC ;AGRAWAL SUDHIR (US); TANG JIN YAN (US); PADMAPRIYA A) 13 October 1994 see claims 1,30; figures 2,3 -----	1,11,42

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 96/04605

Box I Observations where certain claims were found unsearchable (Continuation of item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.:
because they relate to subject matter not required to be searched by this Authority, namely:
Although claims 52-55 (completely and claims 36-51 partly as far as an "in vivo" method is concerned) are directed to a method of treatment of the human/animal body the search has been carried out and based on the alleged effects of the compound/composition.
2. ☐ Claims Nos.:
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 96/04605

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
WO-A-9501985	19-01-95	AU-B- 7258194	06-02-95
GB-A-2225112	23-05-90	DE-A- 3938853	23-05-90
		FR-A- 2639358	25-05-90
		IT-B- 1238025	23-06-93
EP-A-0185494	25-06-86	US-A- 4883750	28-11-89
		DE-A- 3586090	25-06-92
		JP-B- 6044880	15-06-94
		JP-A- 61191964	26-08-86
		US-A- 5521065	28-05-96
		US-A- 5242794	07-09-93
WO-A-9106626	16-05-91	US-A- 5399676	21-03-95
		AU-B- 641219	16-09-93
		AU-B- 7148191	31-05-91
		EP-A- 0539371	05-05-93
		JP-T- 5505101	05-08-93
		US-A- 5527899	18-06-96
WO-A-9417086	04-08-94	NONE	
WO-A-9423028	13-10-94	AU-B- 6527094	24-10-94
		CA-A- 2159350	13-10-94
		EP-A- 0693123	24-01-96